

On the absence of broad Mg II emission-line variability in NGC 3516 during 1996.

M.R. Goad¹, A.P. Koratkar¹, D.J. Axon², K.T. Korista³, and P.T. O’Brien⁴

¹ STScI, 3700 San Martin Drive, Baltimore, MD 21218.

² Division of Physical Sciences, University of Hertfordshire, College Lane, Hatfield, Herts, AL10
9AB, UK

³ Dept. of Physics, Western Michigan University, Kalamazoo, MI 49008.

⁴ Dept. of Physics and Astronomy, University of Leicester, University Road, Leicester, LE1 7RH,
UK.

Abstract

During the 1996 *HST/FOS* monitoring campaign of the Seyfert 1 galaxy NGC 3516, the UV continuum showed a factor of 5 variation; correspondingly the strongest broad UV emission lines (e.g. Ly α , C IV) displayed somewhat weaker, though still significant variations (a factor of 2). In contrast, no variation was detected in the strong broad Mg II + UV Fe II emission-line complex. While historically the flux in the broad Mg II emission line has varied (a factor of 2 within a few years), the profile shape has not been observed to change over the past decade. In contrast, the high ionization lines (HILs) show both emission-line flux and profile changes on relatively short timescales, which appear to be correlated with changes in the continuum level. Significantly, on 1996, February 21, the profile shapes of the HILs (e.g. Ly α , C IV) and low ionization lines (LILs, e.g. Mg II) were identical, at which time the continuum level was at its highest. These results impose stringent constraints upon viable kinematic models of the BLR in NGC 3516, some of which we discuss below.

Subject headings : galaxies : Seyfert — galaxies : individual (NGC 3516) — galaxies :
emission lines — galaxies : kinematics and dynamics

1. Introduction

NGC 3516 is a Seyfert 1 galaxy with a well documented history of short-timescale, large-amplitude continuum and emission-line variability (Koratkar et al. 1996). Historically, *IUE/SWP* spectra of NGC 3516 have also displayed what has previously been referred to as a strong, variable, broad absorption line (VAL, FWHM $\sim 200 \text{ km s}^{-1}$) (Kolman et al. 1993; Walter et al. 1990; Voit et al. 1987), presumed to be similar to that observed in Broad Absorption Line quasars (BALs, Weymann et al. 1991). The subsequent disappearance of the supposed VAL between 1989 and the onset of the 1993 *IUE/SWP* monitoring campaign (Koratkar et al. 1996), may be associated with the increased ionization state of the gas inferred from the observed variation in the strength of the X-ray warm absorber detected by *ROSAT* (Mathur et al. 1997).

However, recent high resolution UV observations of the C IV emission-line region in NGC 3516 with the *HST/GHRS* (Crenshaw et al. 1998), show that NGC 3516 displays multiple intrinsic absorption lines that are all blue-shifted and relatively narrow (FWHM $\sim 500 \text{ km s}^{-1}$), and which appear to remain invariant on timescales of ~ 6 months, and are thought to originate in the narrow-line region gas. In light of these observations, Goad et al. (1999) argue that the supposed VAL is in fact multiple, narrow absorption-line systems which *appear* broad at the lower resolution of IUE. In this picture, the historical variation in the strength of the VAL results from variations in the number of narrow-line clouds coupled with changes in the shape of the underlying broad emission-line.

In an ongoing effort to determine the nature of the broad emission/absorption line regions and their long-term evolution, we obtained 5 HST/FOS cycle 6 observations of NGC 3516, from 1995 December to 1996 November, covering the wavelength range $\lambda\lambda 1150\text{--}3300 \text{ \AA}$. In this paper we present the first results of this study, highlighting the absence of emission-line flux variations in the broad Mg II and UV Fe II emission lines, and its implications in terms of viable kinematic models of the BLR.

1.1. Results from the 1996 HST/FOS monitoring campaign

Goad et al. (1999) showed that during the 1996 HST/FOS monitoring campaign, NGC 3516 displayed the following continuum and emission-line variability characteristics: (i) large amplitude, wavelength dependent, continuum flux variations (a factor of 5 at $\lambda 1365 \text{ \AA}$, c.f. a factor of 3 at $\lambda 2230 \text{ \AA}$), (ii) correlated, large amplitude, UV broad emission-line flux variations (a factor of 2 for both $\text{Ly}\alpha$ and C IV), and (iii) the complete absence of variations in both the strength and shape of the broad Mg II and UV Fe II emission-line complex (Figure 1).

A comparison of the Mg II emission-line strength determined from IUE/LWP observations taken in 1993 (Koratkar et al. 1996) with the present data, show that the flux in Mg II has apparently remained constant for at least 4 years. The absence of significant variability in the Mg II emission-line strength, despite large amplitude continuum and HIL flux variations, has also been noted in lower resolution (pre-1989) IUE/LWP data (Koratkar et al. 1996) when the Mg II emission-line strength was a factor of 2 weaker. Interestingly, the observed doubling of the strength of the broad Mg II emission-line between 1989–1993, appears to coincide with the disappearance of the higher-velocity narrow absorption-line components comprising the so-called VAL. It remains to be seen whether these two phenomena are related. Significantly, despite the notable increase in the Mg II emission-line strength between 1989 and 1993 its shape has remained unchanged for over a decade (Figure 2). Moreover, in the present campaign, the shapes of the broad $\text{Ly}\alpha$, C IV and Mg II emission lines as observed on 1996, February 21, when the continuum was in its *highest* state, are indistinguishable from one another (Figure 3a, N.B. $\text{Ly}\alpha$ is not shown).

2. Implications for kinematic models of the BLR

A viable kinematic model of the BLR in NGC 3516, must explain : (i) the absence of both short-timescale ($\sim$ months) and long-timescale ($\sim$ a few years) variations in the Mg II emission-line flux and shape, despite significant changes in the strength of the ionizing

continuum; and (ii) the remarkable similarity between the line shapes of the HILs and LILs, at high continuum levels, when clearly these lines form under very different physical conditions.

2.1. Flux variability constraints

An absence of response in the Mg II emission line can result for a variety of physical reasons : (i) the continuum band driving the line is invariant; (ii) the emission-line is insensitive to continuum variations; and (iii) the line emitting region is physically extended.

Here we address the shortcomings of each of these scenarios.

(i) While it is true that Mg II and Ly α respond to very different continuum bands (Mg II is mostly sensitive to the continuum from 600–800 eV (Krolik and Kallman 1988) whereas Ly α is driven mainly by Lyman continuum photons), simultaneous multi-wavelength observations of a small number of AGN in the UV, optical and soft X-ray band-passes, indicate that the amplitude of the continuum variations generally increase toward shorter wavelengths (Romano and Peterson 1998). Although we cannot confirm whether such a trend holds at 600–800 eV, the Seyfert 1 galaxies for which simultaneous observations were made in the UV/EUV, NGC 5548 (Marshall et al. 1997) and NGC 4051 (Cagnoni et al. 1998), showed increased variability at shorter wavelengths. Thus, the evidence suggests that the lack of Mg II and Fe II variations is not due to an absence of continuum variability in the bandpass responsible for driving these lines.

(ii) Goad et al. (1993, 1995) and O’Brien et al. (1995) demonstrated that the line responsivity η , the fractional change in line emissivity for a given fractional change in the continuum level, is relatively modest for Mg II ($\eta \sim 0.2 - 0.3$), when compared to the approximately linear response ($\eta \approx 1$) of the HILs. This difference arises because for a single cloud and a fixed continuum shape, the variation in the position of the hydrogen ionization front is generally small compared to the overall extent of the partially ionized

zone where LILs such as Mg II are formed. For a fixed continuum shape this condition holds true over a large range in ionizing continuum luminosity (factors of 10 or more). Although the Mg II line responsivity is small, given the factor of 2 change in the C IV emission-line flux, we should have detected $\sim 40\%$ variation in the Mg II emission-line flux. Since no significant ($<7\%$) variation was detected, low emission-line responsivity cannot be the *sole* explanation for the absence of Mg II emission-line variations.

(iii) Finally, the lack of response in the Mg II emission line can also result if the Mg II line-emitting region is physically large, and has thus yet to respond to the observed continuum variations. While plausible, given the short-timescale response of the C IV emission-line to the continuum variations (~ 4.5 days, Koratkar et al. 1996), it is difficult in this picture to explain the similarity in shape of the C IV and Mg II emission-line profiles at high continuum levels, particularly if the size of the regions where these lines originate differ by a few orders of magnitude.

The similar ionization potential of Fe II and Mg II suggests that these lines are formed in similar regions, hence it is unsurprising that the Fe II emission lines display a similar lack of response to continuum variations (N.B. relatively few of the Fe II lines (although strong) are resonance lines, although others are pumped by the continuum, especially in the presence of extra thermal particle speeds). The smoothness of the Fe II complex in NGC 3516 when compared to similar observations of I Zw I (Goad et al. 1999), a quasar with strong narrow Fe II emission-lines (Laor et al. 1997), supports this finding, and suggests that the Fe II lines in NGC 3516 are indeed produced in high velocity gas ($\text{FWHM} \sim 4500 \text{ km s}^{-1}$).

3. Profile variability constraints

The similarity in shape between the HILs and LILs at the highest continuum levels suggests that in the high-state both C IV and Mg II arise in kinematically similar regions. However, if the lack of response of Mg II is not the result of an absence of variability in the

continuum band driving this line, then the difference in response timescale between the HILs and Mg II suggest that these lines *do* arise in spatially distinct regions, with Mg II formed in a region which extends over several light-months or more. Figure 3b indicates that between the low and high-states, the core of the C IV emission line shows a deficit of response redward of line center from 0 to +4000 km s⁻¹ (even after allowing for possible contamination by the variable absorption-line and narrow emission-line components) and an enhanced response in the far red-wing (> 5000 km s⁻¹). This may result from : (i) reverberation within a radial flow; (ii) a change in the radiation pattern of the ionizing continuum; or (iii) a change in the spatial distribution of the line emitting gas.

Taken together, this evidence places severe constraints upon the spatial distribution and kinematics of the line-emitting gas in NGC 3516. For example, based on this data the following models can be excluded :

- (i) Radial flows at constant velocity, illuminated by either an isotropic or anisotropic continuum source (Goad and Knigge 1999).
- (ii) Radial flows in which Mg II and C IV arise in azimuthally separated cloud populations, either through spatial segregation or anisotropic illumination.
- (iii) A spherical distribution of clouds in Keplerian motion and illuminated by an isotropic continuum source.

We propose two models which may account for the observed emission-line variations: (i) an anisotropic continuum source model and (ii) a terminal wind model (i.e. a flow that accelerates from rest until it reaches constant velocity).

(i) The anisotropic continuum model of Wanders et al. (1995) can broadly reproduce the gross details of the observations reported here. In this model, the continuum is assumed to be comprised of two components, an anisotropic variable component responsible for driving the variations in the HILs and an isotropic non-variable component responsible for producing

the Mg II and Fe II emission. To match the observed variations, the BLR must be comprised of two distinct cloud populations, a low column density population producing significant HIL emission, and a spatially distinct high-column density population producing both HILs and LILs. The variable ionizing continuum component preferentially illuminates the low-column density population, giving rise to the variations in Ly α and C IV emission-lines, whereas the LILs arise predominantly from the high-column density clouds illuminated by the non-variable isotropic continuum component. If the ionization parameter U is large enough, and the EUV is not absorbed, these high column density clouds could also produce significant HIL emission. To account for the similarity in profile shape in the high-state, we further assume that the variable ionizing continuum component becomes more isotropic when brighter, so that the C IV emission arises predominantly from the high-column density clouds. Since the FWHM of the low and high-state C IV emission-line profile is effectively the same, the velocity field cannot be predominantly radial, otherwise a different range in projected velocity for the HILs and LILs will result. Instead a randomized or chaotic flow is favored. While plausible, this model requires a high-degree of fine tuning to produce the detailed profile changes observed in the HILs. Hardest to explain are the reversals in HIL profile shape, from red-asymmetric at low continuum levels to blue-asymmetric at high continuum levels (Figure 3b).

(ii) One possible scenario which may account for the above observations is a terminal wind model for the BLR¹. A terminal wind is the one physical structure which can display the same range in projected velocities on both small and large spatial scales. The wind may be spherical or non-spherical. For a non-spherical wind, for example a bi-conical flow, similar profiles for the HILs and LILs will result *regardless* of their respective radial emissivity distributions, provided that the opening angle of the cone remains constant. However, to

¹The best available variability data on NGC 3516 (Wanders et al. 1993; Koratkar et al. 1996) suggest that the BLR kinematics are not dominated by radial motion, but may be consistent with a mixture of velocity components.

produce profiles similar to that observed here (i.e. broad wings and narrow cores), we require a strong azimuthal dependence on the velocity field, or a significant circularized velocity component to the flow (Goad and Knigge 1999). That is, the simplest terminal flow model in which the velocity field is constant everywhere cannot produce the correct profile shapes. However, hydromagnetic wind models (e.g. Emmering et al. 1992; Bottorff et al. 1997) do exhibit these basic properties. While the emission-line flux variations of the HILs are assumed to be driven by continuum variations, we propose that in this model the changes in profile shape at low continuum levels are in part due to changes in the formation radius of the HILs. Specifically, we propose that in the low-state, conditions within the BLR gas are such that a large fraction of the C IV emission arises at the base of the wind, in the region where the gas begins to accelerate (Goad and Knigge 1999).

4. Conclusions

The picture we envisage is one in which the steady-state profile is dominated by the dynamics of the system, with lines formed in kinematically similar regions, possibly a terminal flow. The stability of the Mg II profile on timescales of several years, provide strong supporting evidence for the existence of such a structure. Moreover, the absence of significant flux variations in the Mg II line on timescales of a few years suggest that this region is physically extended in size. The historic variation in the Mg II emission-line flux, but not its profile, could be the result of an accretion event, a disk instability, or an increase in the wind density and may be linked to the disappearance of the C IV VAL.

Acknowledgements

The results presented here are based upon observations made with the NASA/ESA Hubble Space Telescope, obtained at the Space Telescope Science Institute, which is operated by the Associated Universities for Research in Astronomy, Inc, under NASA contract NAS5-26555.

MRG and AK acknowledge research grant GO-6108 for financial support. We would also like to thank the referee for many helpful suggestions. MRG would also like to thank Christian Knigge for many useful discussions concerning disk-wind modeling.

REFERENCES

- Bottorff, M., Korista, K.T., Shlosman, I., and Blandford, R.D. 1997, ApJ 479, 200.
- Cagnoni, I., Fruscione, A., McHardy, I.M., and Papadakis, I.E. 1998, In : "Dal nano al Tera eV: tutti i colori degli AGN", third italian conference on AGNs, Roma, May 18-21 1998, Memorie S.A.It.
- Crenshaw, D.M., Maran, S.P. and Mushotzky, R.F. 1998, ApJ 496, 797.
- Emmering, R.T., Blandford, R.D., and Shlosman, I. 1992, ApJ 385, 460.
- Goad, M.R., O'Brien, P.T. and Gondhalekar, P.M. 1993, MNRAS 263, 149.
- Goad, M.R. 1995, PhD Thesis, University College London.
- Goad, M.R. et al. 1999, ApJ submitted.
- Goad, M.R. and Knigge, C. 1999, in prep.
- Kolman, M. et al. 1993, ApJ 403, 592.
- Koratkar, A.P. et al. 1996, ApJ 470, 378.
- Krolik, J.H. and Kallman, T.R. 1988, ApJ 324, 714.
- Laor, A., Jannuzi, B.T., Green, R.F., and Boroson, T.A. 1997, ApJ 489, 656.
- Marshall, H.L. et al. 1997, ApJ 479, 222.
- Mathur, S., Wilkes, B.J., and Aldcroft, T. 1997, ApJ 478, 182.
- O'Brien, P.T. Goad, M.R. and Gondhalekar, P.M. 1995, MNRAS 275, 1125.
- Romano, P. and Peterson, B.M. 1998, In : "Structure and Kinematics of Quasar Broad Line Regions", Eds C. M. Gaskell, W. N. Brandt, M. Dietrich, D. Dultzin-Hacyan, and M. Eracleous, ASP Conf. Ser.
- Voit, G.M., Shull, J.M., and Begelman, M.C. 1987, ApJ 316, 573.
- Walter, R., Ulrich, M.-H., Courvoisier, T.J.-L. and Buson, L.M. 1990, A&A 233, 53.
- Wanders, I. et al. 1993, A&A 269, 39.

Wanders, I. et al. 1995, ApJL 453, L87.

Weymann, R.J., Morris, S.L., Foltz, C.B., Hewett, P.C. 1991, ApJ 373, 23.

Figures

Fig. 1.— The high-state (epoch 2, solid line) and low-state (epoch 5, dotted line) continuum subtracted Mg II+Fe II emission-line complex and their difference (dashed line). For clarity, each spectrum has been binned into 2 Å wide bins. The derived errors (poissonian) in the difference spectrum, displaced for clarity, are indicated by the vertical bars. We detect no significant variation in the broad Mg II and UV Fe II emission-line complex. The absorption lines are Galactic in origin.

Fig. 2.— A comparison between the normalized 1996 HST/FOS Mg II emission-line profile (smoothed to the resolution of the IUE/LWP), with that determined from a) the 1993 IUE/LWP monitoring campaign, and b) the IUE/LWP archival data. Note that for the upper panel no scaling between the HST and IUE spectra has been applied, the emission-line fluxes and profile shapes are the same. The solid line in the lower panel shows the old average IUE/LWP Mg II flux scaled relative to the flux measured with HST. While the Mg II emission-line flux clearly varies (a factor of 2.3), the profile shape remains invariant on timescales of several years. The absorption dip blueward of line centre is due to Galactic Mg II.

Fig. 3.— (a) A comparison between the C IV (solid line) and Mg II (dashed line) emission-line profiles as observed on 1996, February 21. Each line has been transformed into velocity space, and normalized to the total flux in a 2000 km s^{-1} wide bin centered at $+2000 \text{ km s}^{-1}$. Note the good agreement between the line profile shapes in the blue and red wings of the lines. The far blue wing of Mg II is contaminated by blended Fe II hence the poor agreement in this region. (b) The high-state (epoch 2, solid line) and low-state (epoch 5, dotted line) continuum subtracted C IV+He II emission lines, scaled to the integrated flux in a 2000 km s^{-1} wide bin centered at -4000 km s^{-1} . The most significant shape changes occur redward of line center, with the core displaying a deficit of response, whereas the far red wing displays an enhanced response. A similar result is found for Ly α .





